

Association vs Causation

Arvid Sjölander

Department of Medical Epidemiology and Biostatistics
Karolinska Institutet

A short course on concepts and methods in Causal Inference

Outline

Association

Causation

Subtle points

Outline

Association

Causation

Preliminaries

- Suppose we are interested in the relation between an exposure, A , and an outcome, Y
- We assume for simplicity that both A and Y are binary
 - We use '0' for 'unexposed/no outcome', and '1' for 'exposed/outcome'
- We assume that population data are available (infinite sample size)
 - No need for p-values, confidence intervals etc
- These conditions are often unrealistic, but are useful for pedagogical purposes
 - Will be relaxed later

Joint probability

- Suppose that the population proportions of A and Y are given by

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

- Among all subjects, 1% are both exposed and have the outcome
- We say that the **joint probability** of $(A = 1, Y = 1)$ is 0.01
- We denote this as $\Pr(A = 1, Y = 1) = 0.01$

Marginal probability

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01
Σ		0.97	0.03

- Among all subjects, 3% have the outcome
- We say that the **marginal probability** of $Y = 1$ is 0.03
- We denote this as $\Pr(Y = 1) = 0.03$

Conditional probability

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

- Among the exposed subjects, $\frac{0.01}{0.01+0.09} = 10\%$ have the outcome
- We say that the **conditional probability** of having the outcome, for exposed subjects, is 0.1
- We denote this as $\Pr(Y = 1|A = 1) = 0.1$

Definition of association and independence

- We say that A and Y are **independent** if the risk of the outcome is the same for exposed and unexposed:

$$\Pr(Y = 1|A = 1) = \Pr(Y = 1|A = 0) = \Pr(Y = 1)$$

- We sometimes write this as $Y \perp\!\!\!\perp A$.
- We say that A and Y are **associated** if the risk of the outcome is different for exposed and unexposed:

$$\Pr(Y = 1|A = 1) \neq \Pr(Y = 1|A = 0) \neq \Pr(Y = 1)$$

- We sometimes write this as $Y \not\perp\!\!\!\perp A$

Example

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

- Are A and Y independent or associated in the table?

Solution

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

$$\Pr(Y = 1|A = 1) = \frac{0.01}{0.01 + 0.09} = 0.1$$

$$\Pr(Y = 1|A = 0) = \frac{0.02}{0.02 + 0.88} = 0.022$$

$$\Pr(Y = 1) = 0.02 + 0.01 = 0.03$$

- $\Pr(Y = 1|A = 1) \neq \Pr(Y = 1|A = 0) \neq \Pr(Y = 1)$, so A and Y are associated

Remark

- There may be several explanations to an association between A and Y
 - A causes Y
 - Y causes A ('reverse causation')
 - A and Y have common causes ('confounding')
- That A and Y are associated only means that certain values of A and Y tend to 'appear together'
 - Why this happens is a different question

Common association measures for binary variables

- The risk difference

$$RD = \Pr(Y = 1|A = 1) - \Pr(Y = 1|A = 0)$$

$$Y \perp\!\!\!\perp A \Leftrightarrow RD = 0$$

- The risk ratio

$$RR = \frac{\Pr(Y = 1|A = 1)}{\Pr(Y = 1|A = 0)}$$

$$Y \perp\!\!\!\perp A \Leftrightarrow RR = 1$$

- The odds ratio

$$OR = \frac{\Pr(Y = 1|A = 1) / \Pr(Y = 0|A = 1)}{\Pr(Y = 1|A = 0) / \Pr(Y = 0|A = 0)}$$

$$Y \perp\!\!\!\perp A \Leftrightarrow OR = 1$$

Example

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

- Compute RD , RR , and OR for the table

Solution

		Y	
		0	1
A	0	0.88	0.02
	1	0.09	0.01

$$\Pr(Y = 1|A = 1) = \frac{0.01}{0.01 + 0.09} = 0.1$$

$$\Pr(Y = 1|A = 0) = \frac{0.02}{0.02 + 0.88} = 0.022$$

$$RD = 0.1 - 0.022 = 0.078$$

$$RR = \frac{0.1}{0.022} = 4.55$$

$$OR = \frac{0.1}{1 - 0.1} / \frac{0.022}{1 - 0.022} = 4.94$$

Conditional association/independence

- Sometimes we wish to stratify data before analysis, e.g:
 - L = 'sex' (0=male, 1=female).
 - $\Pr(Y = 1|A = a, L = 1)$ is the conditional probability of having the outcome, for women with exposure level $A = a$
- **Definition:**
 - A and Y are conditionally independent, given L , if

$$\Pr(Y = 1|A = 1, L) = \Pr(Y = 1|A = 0, L) = \Pr(Y = 1|L)$$

$$Y \perp\!\!\!\perp A \mid L$$

- A and Y are conditionally associated, given L , if

$$\Pr(Y = 1|A = 1, L) \neq \Pr(Y = 1|A = 0, L) \neq \Pr(Y = 1|L)$$

$$Y \not\perp\!\!\!\perp A \mid L$$

Technical note

- In principle, we could have that
 - $\Pr(Y = 1|A = 1, L) = \Pr(Y = 1|A = 0, L)$ for some values of L , and
 - $\Pr(Y = 1|A = 1, L) \neq \Pr(Y = 1|A = 0, L)$ for other values of L
- When we write $Y \perp\!\!\!\perp A \mid L$, we mean that $\Pr(Y = 1|A = 1, L) = \Pr(Y = 1|A = 0, L)$ for **all** values of L
- When we write $Y \not\perp\!\!\!\perp A \mid L$, we mean that $\Pr(Y = 1|A = 1, L) \neq \Pr(Y = 1|A = 0, L)$ for **at least one** value of L

Measures of conditional association

- Conditional risk difference

$$RD(L) = \Pr(Y = 1|A = 1, L) - \Pr(Y = 1|A = 0, L)$$

- Conditional risk ratio

$$RR(L) = \frac{\Pr(Y = 1|A = 1, L)}{\Pr(Y = 1|A = 0, L)}$$

- Conditional odds ratio

$$OR(L) = \frac{\Pr(Y = 1|A = 1, L)}{\Pr(Y = 0|A = 1, L)} / \frac{\Pr(Y = 1|A = 0, L)}{\Pr(Y = 0|A = 0, L)}$$

Causal models

- The sufficient-component cause model (Rothman)
- Potential outcomes, counterfactuals (Rubin, Robins)
- Structural equations, causal diagrams (Pearl)

Outline

Association

Causation

Subtle points

Relation between models

- All common causal models are essentially equivalent, from a mathematical perspective
 - different languages, same content
- To define 'causation', we will mostly rely on the potential outcome model, but borrow from the other models as well

Motivating example

- August has been smoking 5 cigs/day since he was 15 years old. At the age of 60 he develops liver cancer
- *Did the smoking cause the cancer?*

Human reasoning about cause and effects

- We mentally compare two scenarios:
 - the outcome when the exposure is present
 - the outcome when the exposure is absent
- **everything else equal**
- If the two outcomes differ, then we say that the exposure has a causal effect
 - causative or preventative

Ideal data

- Let Y_a be the outcome that we would observe, for a given subject, if the subject potentially received exposure level a
 - Y_1 is the outcome under exposure
 - Y_0 is the outcome under non-exposure
- Y_1 and Y_0 are referred to as **potential outcomes**
- Ideally - **and very unrealistically** - we could observe both potential outcomes for any given subject

subject	Y_1	Y_0
August	1	0
Selma	0	0
Fjodor	1	1

Subject-specific causal effects

subject	Y_1	Y_0
August	1	0
Selma	0	0
Fjodor	1	1

- A has a causal effect on Y , for a given subject, if the potential outcomes Y_1 and Y_0 differ for this subject
 - For August, the exposure has an effect: $Y_1 \neq Y_0$
 - For Selma and Fjodor, the exposure has not effect; $Y_1 = Y_0$

Observed data

- August is exposed ($A = 1$). Thus, for August
 - Y_1 is observed and equal to the factual outcome Y
 - Y_0 is unobserved, or **counterfactual**
- Selma and Fjodor are unexposed ($A = 0$). Thus, for Selma and Fjodor
 - Y_0 is observed and equal to the factual outcome Y
 - Y_1 is unobserved, or **counterfactual**

subject	A	Y	Y_1	Y_0
August	1	1	1	?
Selma	0	0	?	0
Fjodor	0	1	?	1

A fundamental problem of causation



- It is very difficult to say whether the exposure causes the outcome for a specific subject
 - because we cannot observe the same subject under two exposure levels simultaneously

Population effects

- Fortunately, it is much easier to make causal claims on population levels
 - e.g. 'if everybody would quit smoking, then the incidence of liver cancer would decrease by 15%'
 - more later

Population causal effects

- $\Pr(Y_a = 1)$ is the proportion of subjects that would develop the outcome, if **everybody** would receive exposure level a
 - The probability of the outcome if everybody would receive a
- A has a population causal effect on Y if

$$\Pr(Y_1 = 1) \neq \Pr(Y_0 = 1)$$

- A has no population causal effect on Y if


$$\Pr(Y_1 = 1) = \Pr(Y_0 = 1)$$

Technical note


- In statistics, we use
 - upper case letters (e.g. A , Y) for random variables
 - lower case letters (e.g. a , y) for fixed numbers
- When writing Y_a , we consider the exposure to be fixed to a (0 or 1)
- When writing $\Pr(Y_a = 1)$, we consider a scenario where the exposure is fixed to a for everybody

Association vs Causation

- Association:



- Causation:



Measures of causal effects for binary variables

- The causal risk difference

$$CRD = \Pr(Y_1 = 1) - \Pr(Y_0 = 1)$$

Causal null hypothesis $\Leftrightarrow CRD = 0$

- The causal risk ratio

$$CRR = \frac{\Pr(Y_1 = 1)}{\Pr(Y_0 = 1)}$$

Causal null hypothesis $\Leftrightarrow CRR = 1$

- The causal odds ratio

$$COR = \frac{\Pr(Y_1 = 1) / \Pr(Y_1 = 0)}{\Pr(Y_0 = 1) / \Pr(Y_0 = 0)}$$

Causal null hypothesis $\Leftrightarrow COR = 1$

Example

subject	Y_1	Y_0
1	0	0
2	1	0
3	0	0
4	1	1
5	0	0
6	1	1
7	1	1
8	1	1
9	0	0
10	1	0

- Compute CRD , CRR , and COR

Solution

subject	Y_1	Y_0
1	0	0
2	1	0
3	0	0
4	1	1
5	0	0
6	1	1
7	1	1
8	1	1
9	0	0
10	1	0

$$\Pr(Y_1 = 1) = 6/10 = 0.6$$

$$\Pr(Y_0 = 1) = 4/10 = 0.4$$

$$CRD = 0.6 - 0.4 = 0.2$$

$$CRR = \frac{0.6}{0.4} = 1.5$$

$$COR = \frac{0.6}{1 - 0.6} / \frac{0.4}{1 - 0.4} = 2.25$$

Conditional causal effects

- Conditional causal risk difference

$$CRD(L) = \Pr(Y_1 = 1|L) - \Pr(Y_0 = 1|L)$$

- Conditional causal risk ratio

$$CRR(L) = \frac{\Pr(Y_1 = 1|L)}{\Pr(Y_0 = 1|L)}$$

- Conditional causal odds ratio

$$COR(L) = \frac{\Pr(Y_1 = 1|L)}{\Pr(Y_1 = 0|L)} / \frac{\Pr(Y_0 = 1|L)}{\Pr(Y_0 = 0|L)}$$

A brief remark

- We have seen that both association and causation can be quantified with risk differences, risk ratios, and odds ratios
- For convenience, we will mostly focus on risk ratios
- Everything that we say holds for risk differences and odds ratios as well

Outline

Association

Causation

Subtle points

When is a counterfactual well defined?

- 'Well defined' = we have a clear understanding of what the counterfactual represents 'in real life'
- Are all counterfactuals well defined?
- If some counterfactuals are not well defined, then causal effects based on these are not well defined either

Example

- Define $A = 1$ if $\text{BMI} > 30$, and $A = 0$ if $\text{BMI} < 30$
- Certain diseases occur more frequently in obese than in non-obese, i.e.

$$\Pr(Y = 1 | A = 1) > \Pr(Y = 1 | A = 0)$$

- Does 'obesity' have a causal effect on the risk for disease?

$$\Pr(Y_1 = 1) \neq \Pr(Y_0 = 1)?$$

Quite a vague question

- Translated into plain English, the counterfactual comparison reads
 - 'what would the risk be if everybody had $\text{BMI} > 30$ compared to if everybody had $\text{BMI} < 30$ '?
- But what does 'if everybody had $\text{BMI} > 30$ ' really mean?
 - everybody short or heavy?
 - everybody fat or muscular?
 - everybody belly fat or hips fat?
- The outcome is probably very different under these alternative counterfactual scenarios
 - Unless we specify more precisely what scenario we refer to, the counterfactual outcome is not well defined

An important difference between association and causation

- In order for the causal effect of A on Y to be well defined we require that
 - we can tell whether an observed subject has $A = 1$ or $A = 0$
 - we agree on what it means that an observed subject with $A = 0$ **would have had** $A = 1$, and vice versa
- In order for the association between A and Y to be well defined, only the first condition is required
 - Because the concept of association is only based on factual observations, not on counterfactuals

‘Some counterfactuals are ill-defined, most are somewhat vague, but many are useful’

Lewis, 1973

Summary

- Association is not equal to causation
- To define causation, we use **potential outcomes** and **counterfactuals**
- Not all counterfactuals (and causal effects) are well defined